

Accepted Manuscript

Boiling factors in the pot of type 1 diabetes mellitus management: the role of exercise

Roberto Codella

PII: S0306-9877(17)30585-6


DOI: <http://dx.doi.org/10.1016/j.mehy.2017.06.026>

Reference: YMEHY 8607

To appear in: *Medical Hypotheses*

Received Date: 4 June 2017

Accepted Date: 28 June 2017



Please cite this article as: R. Codella, Boiling factors in the pot of type 1 diabetes mellitus management: the role of exercise, *Medical Hypotheses* (2017), doi: <http://dx.doi.org/10.1016/j.mehy.2017.06.026>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

- Correspondence -

**Boiling factors in the pot of type 1 diabetes mellitus management:
the role of exercise**

Roberto Codella

Department of Biomedical Sciences for Health, Università degli Studi di Milano, Milan, Italy
Metabolism Research Center, IRCCS Policlinico San Donato, San Donato Milanese, Italy

Letter counts:

| | |
|------------|-------------|
| Title | 15 (words) |
| Body text | 363 (words) |
| References | 11 |

Corresponding author

Roberto Codella, Ph.D.
Department of Biomedical Sciences for Health
Università degli Studi di Milano
Via Fratelli Cervi 93, 20090 Segrate (Milano)
Phone: +39 02 50330300
Fax: +39 02 50315152
E-mail: roberto.codella@unimi.it

Dear Editor,

I found the article by Farinha et al. titled “Exercise for type 1 diabetes mellitus management: General considerations and new directions” [1] of much professional and personal interest. The authors overviewed the benefits of training subjects with type 1 diabetes mellitus (T1D), particularly focusing on modality of anaerobic - high intensity exercise - for reducing exercise-induced hypoglycemic episodes. They ultimately hypothesized a training protocol combining strength training and high intensity interval training, capable to minimize exercise-associated drops in glucose levels in T1D. This speculation makes two complementary observations that can be further discussed.

First, the authors pointed at the crucial effects of exercise in mitigating oxidative stress and inflammation, which are potentially triggering the autoimmune process towards the insulin secreting cells. Such background information, fascinatingly challenged by the same authors in previous studies [2,3], introduces another effect exerted by exercise: a putative immunomodulation of systemic functions to both T1D and inflammation [4]. Not only exercise would elicit glucose-lowering effects [5] but also, according to our epidemiological screening, it would be able to prolong the “honeymoon” phase in active people, like athletes with T1D [6].

Second, if the training orchestrated by the Authors were likely to stem metabolic destabilization post-effort (as several recent data are similarly suggesting [7–9]) then the program would hardly cover, in the long-term, all

the supposed health-related adaptations for subjects with T1D without including a decent contribution from aerobic exercise. Although the optimal glucose control remains of paramount relevance in T1D, conclusively, long-term health and life expectancy are still favoured by regular aerobic exercise. Even in islet-transplanted ultra-marathon runners with T1D, we ascertained that an abundant volume of aerobic training counteracted diabetic symptoms and alleviated the side effects of immunosuppressive drugs, over a 10-year follow up [10].

Besides, an exercise program based solely on high intensity mode may not be easily administrated to everybody, without a certain degree of familiarization. Nowadays, multiple technologies are available and can facilitate an adequate training, under this perspective [11].

Certainly, all these factors, albeit conceptually simple, are boiling in the same pot and they need to be carefully managed in order to prepare personalized recipes benefiting countless people living with T1D.

References

- [1] Farinha JB, da Silva Krause M, Rodrigues-Krause J, Reischak-Oliveira A, Kahn BB, Youn JH. Exercise for type 1 diabetes mellitus management: General considerations and new directions. *Med Hypotheses* 2017;37:139–46.
doi:10.1016/j.mehy.2017.05.033.

- [2] da Silva Krause M, de Bittencourt PIH. Type 1 diabetes: can exercise impair the autoimmune event? The L-arginine/glutamine coupling hypothesis. *Cell Biochem Funct* 2008;26:406–33. doi:10.1002/cbf.1470.
- [3] da Silva Krause M, Bittencourt A, Homem de Bittencourt PI, McClenaghan NH, Flatt PR, Murphy C, et al. Physiological concentrations of interleukin-6 directly promote insulin secretion, signal transduction, nitric oxide release, and redox status in a clonal pancreatic β -cell line and mouse islets. *J Endocrinol* 2012;214:301–11. doi:10.1530/JOE-12-0223.
- [4] Codella R, Terruzzi I, Luzi L. Why should people with type 1 diabetes exercise regularly? *Acta Diabetol* 2017. doi:10.1007/s00592-017-0978-x.
- [5] Codella R, Lanzoni G, Zoso A, Caumo A, Montesano A, Terruzzi IM, et al. Moderate Intensity Training Impact on the Inflammatory Status and Glycemic Profiles in NOD Mice n.d. doi:10.1155/2015/737586.
- [6] Codella R, Luzi L, Inverardi L, Ricordi C. The anti-inflammatory effects of exercise in the syndromic thread of diabetes and autoimmunity. *Eur Rev Med Pharmacol Sci* 2015;19:3709–22.
- [7] Steckling F, Farinha J, Santos D, Bresciani G, Mortari J, Stefanello S, et al. High Intensity Interval Training Reduces the Levels of Serum Inflammatory Cytokine on Women with Metabolic Syndrome. *Exp Clin Endocrinol Diabetes* 2016;124:597–601. doi:10.1055/s-0042-111044.
- [8] Bally L, Zueger T, Buehler T, Dokumaci AS, Speck C, Pasi N, et al. Metabolic and hormonal response to intermittent high-intensity and continuous moderate intensity exercise in individuals with type 1 diabetes: a randomised crossover study. *Diabetologia* 2016;59:776–84. doi:10.1007/s00125-015-3854-7.
- [9] Cassidy S, Thoma C, Houghton D, Trenell MI. High-intensity interval training: a review of its impact on glucose control and cardiometabolic health. *Diabetologia* 2017;60:7–23. doi:10.1007/s00125-016-4106-1.
- [10] Codella R, Adamo M, Maffi P, Piemonti L, Secchi A, Luzi L. Ultra-marathon 100 km in an islet-transplanted runner. *Acta Diabetol* 2016;1–4. doi:10.1007/s00592-016-0938-x.
- [11] Adamo M, Codella R, Casiraghi F, Ferrulli A, Macrì C, Bazzigaluppi E, et al. Active

subjects with autoimmune type 1 diabetes have better metabolic profiles than sedentary controls. Cell Transplant 2017;26:23–32(10).

ACCEPTED MANUSCRIPT

Conflict of Interest

The author declares that he has no conflict of interest.

Funding and Acknowledgements

This article-letter did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

ACCEPTED MANUSCRIPT